

# The National Immunisation Schedule

Age	Disease to protect against	Vaccine
Pregnancy	Tetanus + diphtheria + whooping cough (pertussis)	<b>Boostrix<sup>®</sup></b>
	Influenza	Brand varies.
6 Weeks	Rotavirus (first dose must be given before 15 weeks)	<b>Rotarix<sup>®</sup></b> (oral)
	Diphtheria + tetanus + whooping cough (pertussis) + polio + hepatitis B + <i>Haemophilus influenzae</i> type b (Hib)	<b>Infanrix<sup>®</sup> hexa</b>
	Pneumococcal disease	<b>Synflorix<sup>®</sup></b>
3 Months	Rotavirus (second dose must be given before 25 weeks)	<b>Rotarix<sup>®</sup></b> (oral)
	Diphtheria + tetanus + whooping cough + polio + hepatitis B + <i>Haemophilus influenzae</i> type b (Hib)	<b>Infanrix<sup>®</sup> hexa</b>
5 Months	Diphtheria + tetanus + whooping cough + polio + hepatitis B + <i>Haemophilus influenzae</i> type b (Hib)	<b>Infanrix<sup>®</sup> hexa</b>
	Pneumococcal disease	<b>Synflorix<sup>®</sup></b>
12 Months	Measles + mumps + rubella	<b>Priorix<sup>®</sup></b>
	Pneumococcal disease	<b>Synflorix<sup>®</sup></b>
15 Months	<i>Haemophilus influenzae</i> type b (Hib)	<b>Hiberix<sup>®</sup></b>
	Measles + mumps + rubella	<b>Priorix<sup>®</sup></b>
	Chickenpox (varicella)	<b>Varivax<sup>®</sup></b>
4 Years	Diphtheria + tetanus + whooping cough + polio	<b>Infanrix<sup>®</sup> IPV</b>
11-12 Years	Tetanus + diphtheria + whooping cough	<b>Boostrix<sup>®</sup></b>
	Human papillomavirus (HPV)	<b>Gardasil<sup>®</sup> 9</b> (2 doses, 6 months apart)
45 Years	Tetanus + diphtheria + whooping cough	<b>Boostrix<sup>®</sup></b>
65 Years	Tetanus + diphtheria + whooping cough	<b>Boostrix<sup>®</sup></b>
	Influenza	Brand varies.
	Shingles	<b>Zostavax<sup>®</sup></b>